

Thuricide®

Bacillus Thuringiensis (BT)

C · o · n · c · e · n · t · r · a · t · e

ACTIVE INGREDIENTS:

*Bacillus thuringiensis Kurstaki, potency of 4,000 I.U. (at least 6 million viable spores per milligram)	0.80%
OTHER INGREDIENTS:	99.20%
TOTAL	100.00%

There is no direct relationship between intended (potency) and the Percent Active Ingredient by Weight.

Contains petroleum hydrocarbon solvent
Equivalent to 4.0 billion I.U. per quart THURICIDE® - HPC:
Reg. Trademark of Sandoz, Inc.

EPA Est. No. 4-NY-1

EPA Reg. No. 4-226

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

(See Back Panel for Additional Precautionary Statements)



**Controls
moth larvae
(caterpillars),
certain leaf
eating worms
and Gypsy
Moths.**

Trusted since 1926
BONIDE PRODUCTS, INC.

Complete directions for use have been removed!

KEEP OUT OF REACH OF CHILDREN

Dispose of properly

Buyers Guarantee limited to label claims

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Questions, Concerns or Comments?

Write to

BONIDE PRODUCTS, INC.

Consumer Services

6301 Sutliff Road

Oriskany, NY 13424

Lift & Peel Here →

**BACILLUS THURINGIENSIS (BT)
THURICIDE® Concentrate**

CONTROLS alfalfa and redhumped caterpillars, cabbage looper, omnivorous and fruit tree leaf roller...**USE** on alfalfa, broccoli, cabbage, mustard greens, melons, tomatoes and others as listed...**MIX** 1 to 4 tsp. per gallon of water. **Mixture ratios vary. Read and use according to label directions. APPLY** using hose or pressurized sprayers. Can be applied up to the day of harvest.

Store and transport in an upright position.
Buyers Guarantee Limited to Label Claims.
EPA Est. No. 4-NY-1 EPA Reg. No. 4-226
Manufactured by
BONIDE PRODUCTS, INC.
6301 Suttiff Road
Oriskany, NY 13424

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION - Bonide (BT) is a bacterium which is selectively toxic to many moth and butterfly larvae (Caterpillars). The insects stop feeding and die within 2-3 days of ingestion. There is no residue problem, and being exempt from tolerance requirements, this product may be applied at recommended amounts up to the day of harvest.

Apply at first sign of infestation, and repeat at weekly intervals as needed to maintain control. Bonide BT may be applied by hose or pressurized sprayers. Apply thoroughly to upper and lower leaf surfaces, but not to the point of excessive run-off.

Page: One

Mixing: Shake or stir thoroughly before use. Add recommended amount of product to water in jar or tank while shaking or stirring. A sticker and evaporation retardant (such as molasses) may be added to the spray mix. Do not store diluted spray mix for more than 12 hours.

Gypsy Moth - To use on trees infested with gypsy moth caterpillars, apply when caterpillars are small (2nd or 3rd instars), and repeat treatment in two weeks. Use at the rate of 4 teaspoons/gal. water.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Mustard Greens, Turnip Greens-to control cabbage looper use at a rate of 2-4 tsp. per gal. of water. To control imported cabbage worm use at a rate of 1-3 tsp. per gal. of water.

Page: Two

Beans, Celery, Lettuce, Spinach-to control cabbage looper use at a rate of 2-4 tsp. per gal. of water.

Melons, Potatoes, Cucumbers (Southeast only)-to control cabbage looper use at a rate of 1-3 tsp. per gal. of water.

Tobacco-to control cabbage looper use at a rate of 2-3 tsp. per gal. of water. To control tobacco budworm, tobacco hornworm use at a rate of 1-2 tsp. per gal. of water.

Tomatoes-to control cabbage looper use at a rate of 1-3 tsp. per gal. of water. To control tomato hornworm use at a rate of 1-2 tsp. per gal. of water.

Page: Three

Page: Four

Watermelons (Florida)-to control rindworm complex use at a rate of 1-2 tsp. per gal. of water.

Grapes-to control grape leaf folder use at a rate of 2-4 tsp. per gal. of water. To control omnivorous leaf roller (California) use at a rate of 1-3 tsp. per gal. of water.

Oranges (Ground application only - Arizona and California)-to control fruit tree leaf roller use at a rate of 2-4 tsp. per gal. of water. To control orange dog use at a rate of 1-1 1/2 tsp. per gal. of water.

Walnuts (Ground application only - California)-to control redhumped caterpillar use at a rate of 1/2-2 tsp. per gal. of water.

STORAGE AND DISPOSAL

Store in a cool, dry place, and in an area inaccessible to children or pets. Store and transport in upright position. Securely wrap container in newspaper and place in trash collection.

PRECAUTIONARY STATEMENTS

CAUTION

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams or ponds. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

Page: Five

Recommendations for use of this product are based upon tests believed to be reliable. Buyer assumes all risk for use of this product when not used in strict accordance with label directions, or when used under abnormal conditions not reasonably foreseeable to the seller.

For information on pesticide products (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.

Page: Six

Page: Seven